

# Vision Problems in Ohio

Prevalence of Adult Vision Impairment and  
Age-Related Eye Disease in Ohio



Published by:



1500 W. Third Ave, Ste. 200, Columbus, OH 43212  
614-464-2020 800-301-2020 Fax: 614-481-9670

[preventblindnessohio@compuserve.com](mailto:preventblindnessohio@compuserve.com)

[www.preventblindness.org](http://www.preventblindness.org)

*Source: Vision Problems in the U.S., a publication of Prevent Blindness America in cooperation with the National Eye Institute, National Institutes of Health, U.S. Dept. of Health and Human Services. Published 2002.*

# Prevalence of Adult Vision Impairment and Age-Related Eye Disease in Belmont County (70,226)

## Vision Loss Among Older Ohioans Is Increasing

More Americans than ever are facing the threat of blindness from age-related eye disease. Over 187,000 Ohioans age 40 and older are currently legally blind or visually impaired, largely resulting from the eye diseases of diabetic retinopathy, cataract, glaucoma, and age-macular degeneration (AMD). The number of older Americans affected by these diseases is expected to double over the next 30 years as the Baby Boomer generation ages. **More than 2.5 million Ohioans will be affected.** Unfortunately blindness and vision impairment represent a significant burden, not only to those affected by sight loss, but to our Ohio economy as well. It is estimated that blindness and vision impairment cost Ohio \$160,000,000 annually in benefits and lost taxable income.

Below are statistics for adult vision impairment and age-related eye disease, including:

- **Visual Impairment** is defined as having 20/40 or worse in the better eye, even with eyeglasses.
- **Blindness** is defined as visual acuity with best correction in the better eye worse than or equal to 20/200 or a visual field extent of less than 20 degrees in diameter.
- **Myopia** is the same as nearsightedness.
- **Age-Related Macular Degeneration** creates a loss of sharp, central vision. No successful treatment is currently available.
- **Cataract** is a clouding of the eye's naturally clear lens. This is usually treatable.
- **Diabetic Retinopathy** involves abnormal growth of blood vessels in the back of the eye which can leak. Management of diabetes and regular preventative eye exams are the best defense against developing diabetic retinopathy.
- **Glaucoma** causes the loss of peripheral or side vision. Once vision is lost, it cannot be restored; however, further loss of remaining vision can usually be prevented with treatment.

	Total	Male	Female	White	Black	Hispanic	Other
<b>Visual Impairment, including Blindness (est. cases): Age 40+</b>	1,041	335	706	1,019	20	2	*
<b>Blindness (est. cases): Age 40+</b>	321	96	225	312	8	*	*
<b>Myopia: Age 40+</b>	9,868	4,373	5,495	9,726	122	20	*
<b>Age-related Macular Degeneration: Age 50+</b>	554	189	355	548	6	1	*
<b>Cataract: Age 40+</b>	6,439	2,376	4,063	6,314	112	12	*
<b>Diabetic Retinopathy: Age 18+</b>	1,403	658	745	1,343	52	5	4
<b>Glaucoma: Age 40+</b>	605	204	401	574	29	2	*

*Source: Vision Problems in the U.S., a publication of Prevent Blindness America in cooperation with the National Eye Institute, National Institutes of Health, U.S. Dept. of Health and Human Services. Published 2002. Visit [www.preventblindness.org](http://www.preventblindness.org).*

Data Provided By:



1500 W. Third Avenue, Suite 200 , Columbus, OH 43212  
 614-464-2020 800-301-2020 Fax: 614-481-9670  
[preventblindnessohio@compuserve.com](mailto:preventblindnessohio@compuserve.com)  
[www.preventblindness.org](http://www.preventblindness.org)